

**TITLE:**

**Mobile PEAR transcription factors integrate positional cues to prime cambial growth**

**AUTHORS:**

**Shunsuke Miyashima<sup>2,3\*</sup>, Pawel Roszak<sup>1,2\*</sup>, Iris Sevilem<sup>2\*</sup>, Koichi Toyokura<sup>1,7\*\*</sup>, Bernhard Blob<sup>1\*\*</sup>, Jung-ok Heo<sup>1,2</sup>, Nathan Mellor<sup>8</sup>, Hanna Help-Rinta-Rahko<sup>2</sup>, Sofia Otero<sup>1</sup>, Wouter Smet<sup>4,5,6</sup>, Mark Boekschoten<sup>14</sup>, Guido Hooiveld<sup>14</sup>, Kayo Hashimoto<sup>3,9</sup>, Ondřej Smetana<sup>2</sup>, Riccardo Siligato<sup>2</sup>, Eva-Sophie Wallner<sup>12</sup>, Ari Pekka Mähönen<sup>2</sup>, Yuki Kondo<sup>10</sup>, Charles W Melnyk<sup>1,11</sup>, Thomas Greb<sup>12</sup>, Keiji Nakajima<sup>3</sup>, Rosangela Sozzani<sup>13</sup>, Anthony Bishopp<sup>8</sup>, Bert De Rybel<sup>4, 5, 6 #</sup>, Ykä Helariutta<sup>1, 2 #</sup>**

**AFFILIATION:**

1: The Sainsbury Laboratory, University of Cambridge, Cambridge, CB2 1LR, UK

2: Institute of Biotechnology, HiLIFE/Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, and Viikki Plant Science Centre, University of Helsinki, FIN-00014, Finland

3: Graduate School of Science and Technology, Nara Institute of Science and Technology, Nara, 630-0192 Japan

4: Ghent University, Department of Plant Biotechnology and Bioinformatics, 9052 Ghent, Belgium

5: VIB Center for Plant Systems Biology, 9052 Ghent, Belgium

6: Wageningen University, Laboratory of Biochemistry, 6708 WE, Wageningen, The Netherlands

7: Department of Biological Sciences, Graduate School of Science, Osaka University, Toyonaka, Osaka, 560-0043, Japan

8: Centre for Plant Integrative Biology (CPIB) and School of Biosciences, University of Nottingham, Nottingham, LE12 5RD, UK

9: Graduate School of Humanities and Sciences, Nara Women's University, Nara, 630-8506 Japan

10: Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Tokyo, 113-0033, Japan

11: Department of Plant Biology, Swedish University of Agricultural Sciences, 756 51 Uppsala, Sweden

12: Centre for Organismal Studies (COS), Heidelberg University, 69120 Heidelberg, Germany

13: Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC 27695, United States.

14: Nutrition, Metabolism and Genomics Group, Division of Human Nutrition, Wageningen University, 6708 WE Wageningen, The Netherlands

# Corresponding author. Email: Ykä Helariutta: yrjo.helariutta@slcu.cam.ac.uk; Bert De Rybel: beryb@psb.vib-ugent.be

\* These authors contributed equally to this work.

\*\* These authors contributed equally to this work.

While apical growth in plants initiates upon seed germination, radial growth is only primed during early ontogenesis in procambium cells and activated later by the vascular cambium<sup>1</sup>. Although it is not known how radial growth is organized and regulated in plants, this system resembles the developmental competence observed in some animal systems, in which pre-existing patterns of developmental potential are established early on<sup>2,3</sup>. Here we show that the initiation of radial growth occurs around early protophloem sieve element (PSE) cell files of the root procambial tissue in *Arabidopsis*. In this domain cytokinin signalling promotes expression of a pair of novel mobile transcription factors, PHLOEM EARLY DOF (PEAR1, PEAR2) and their four homologs (DOF6, TMO6, OBP2 and HCA2), collectively called PEAR proteins. The PEAR proteins form a short-range concentration gradient peaking at PSE and activating gene expression that promotes radial growth. The expression and function of PEAR proteins are antagonized by well-established polarity transcription factors, HD-ZIP III<sup>4</sup>, whose expression is concentrated in the more internal domain of radially non-dividing procambial cells by the function of auxin and mobile miR165/166. The PEAR proteins locally promote transcription of their inhibitory *HD-ZIP III* genes, thereby establishing a negative feedback loop that forms a robust boundary demarking the zone of cell divisions. Taken together, we have established a network, in which the PEAR - HD-ZIP III module integrates spatial information of the hormonal domains and miRNA gradients during root procambial development, to provide adjacent zones of dividing and more quiescent cells as a foundation for further radial growth.

Cambial growth in plants is initiated within the procambial tissues of the apical meristems through periclinal (i.e. longitudinal) divisions associated with formation of the vascular tissues xylem and phloem<sup>1</sup> (Extended Data Fig. 1a). It has been established that during procambial development in *Arabidopsis* roots there are distinct domains for high auxin and cytokinin signalling, which mark the regions for further development of xylem and phloem/procambium, respectively<sup>5-8</sup>. To accurately map the spatial distribution of the periclinal divisions, we established a new nomenclature for the root procambial cells, including PSE-lateral neighbours (PSE-LN) as cells directly contacting both PSE and the pericycle, the outer procambial cells (OPC) as procambial cells adjacent to the pericycle but not contacting PSE, and SE-internal neighbours (PSE-IN) as cells located internal to and directly contacting PSE (Fig. 1a). Both the PSE cell and PSE-LN showed higher activity of periclinal cell division than the OPC and PSE-IN (Fig. 1b, Extended Data Fig. 1b-d and Supplementary Information). We observed virtually no periclinal divisions in metaxylem (MX) and internal procambial cells (IPC) (Fig. 1b). Furthermore, blocking symplastic transport genetically<sup>9</sup> between the PSE and the surrounding cells results in a dramatic reduction in the number of cell files, not only in PSE lineage but also in the PSE-LN lineage (Extended Data Fig. 2a-e). Thus, the proliferative activity in procambium is centred on and around PSE and may involve symplastic intercellular signals.

By searching *in silico* for transcription factors enriched in early PSE<sup>10</sup> (Extended Data Fig. 3a), we identified a pair of DOF transcription factors<sup>11</sup>, *PHLOEM-EARLY-DOF 1* (*PEAR1*)/*DOF2.4* and *PEAR2/DOF5.1*<sup>12</sup> (Extended Data Fig. 3b). RNA *in situ* hybridization and transcriptional fusion constructs validated that both *PEAR1* and *PEAR2* are transcribed specifically in PSE cells (Fig. 1c and Extended Data Fig. 3d). However, fluorescent tagged versions of the PEAR proteins show localization not only in PSE but also in PSE neighbouring cells (PSE-LN and PSE-IN), indicating that these proteins move across short ranges via plasmodesmata (Fig. 1d, Extended Data Fig. 2f-g, 3d and 4a-d).

83 We next investigated whether the loss-of-function of these genes would lead to a phenotype  
 84 corresponding to the one observed when symplastic transport is compromised (Extended Data Fig 2c).  
 85 However, we did not find such phenotype in single or double mutants corresponding to *PEAR1* and  
 86 *PEAR2* (Extended Data Fig. 5a and b). We subsequently identified *DOF1.1/OBP2*<sup>13</sup>, *DOF3.2/DOF6*<sup>14</sup>,  
 87 *DOF5.6/HCA2*<sup>15</sup>, and *DOF5.3/TMO6*<sup>16</sup> as additional PSE specific/abundantly expressed *DOF* genes  
 88 with a broader gene product localization (Extended data Fig 3d). Furthermore, overexpression of any  
 89 of these six loci results in an increased number of cell files (Extended Data Fig. 3c). In addition, we  
 90 observed that *DOF6*, *HCA2* and *TMO6* are upregulated in *pear1 pear2* double mutant apparently as a  
 91 compensation response (Extended Data Fig. 3e, see also Supplementary Information). Among the  
 92 several higher order combinatorial mutants involving all six genes, we found the *pear1 pear2 tmo6*  
 93 triple mutant to display reduced radial growth variably (Fig. 2a, c and f), while the corresponding three  
 94 double mutants did not show this phenotype (Extended Data Fig. 5b). Furthermore, the *pear1 pear2*  
 95 *dof6 tmo6* quadruple mutant results in all plants with further, uniformly reduced radial growth  
 96 corresponding to the line with compromised symplastic trafficking (Fig. 2d, f and Extended Data Fig.  
 97 2c), indicating that these four mobile PEAR proteins play a major role in radial growth. In addition,  
 98 the *pear1 pear2 dof6 obp2 hca2* quintuple mutant resulted in a population of slowly elongating roots  
 99 (around 30 per cent, n=300) with a reduction in radial growth (Fig. 2b and f), whereas the  
 100 corresponding five quadruple mutants for the five genes did not display a strong phenotype (Extended  
 101 Data Fig. 5b). The introduction of *obp2* and *hca2* mutations into the *pear1 pear2 dof6 tmo6* quadruple  
 102 background (resulting in the *pear1 pear2 dof6 tmo6 obp2 hca2* hextuple mutant) did not result in  
 103 further reduced radial growth (Fig. 2d-f), collectively suggesting a significant but minor contribution  
 104 of *OBP2* and *HCA2*. We were able to suppress the phenotype of the quintuple and/or hextuple mutants  
 105 with all six genes (Extended Data Fig. 5c, d and see Supplementary Information). Collectively these  
 106 data indicate that the mobile PEAR proteins redundantly control cell proliferation in and around PSE  
 107 cells. Their effects are likely to be both cell autonomous and/or non-cell autonomous as several  
 108 putatively direct target genes, including a central regulator of phloem formation *SUPPRESSOR OF*  
 109 *MAX2 1-LIKE3 (SMXL3)*<sup>17</sup>, are expressed in both PSE and its surrounding cells (Fig. 2g-h and  
 110 Extended Data Fig. 6, also see Supplementary Information). Moreover, ectopic expression of *SMXL3*  
 111 is sufficient to enhance periclinal cell divisions (Extended Data Fig. 6j).

112 Earlier studies have highlighted cytokinins in regulating procambial cell proliferation<sup>6,8</sup>.  
 113 During root development, cytokinin signalling reporter, *pARR5::RFPer*<sup>18</sup> is initially activated and  
 114 maintained in PSE and its surrounding procambial cells, later becoming concentrated in the procambial  
 115 cells neighbouring to the xylem cells, while auxin response is maintained in xylem domain<sup>7,8</sup>  
 116 (Extended Data Fig. 7a, and see Supplementary Information). Cytokinin signalling reporter partially  
 117 overlaps with the *PEAR1* transcriptional domain (Fig. 3a). Exogenous cytokinin application rapidly  
 118 increased the level of some of the PEAR transcripts (Extended Data Fig. 7b), and sustained cytokinin  
 119 treatment resulted in a radial expansion of *PEAR* expression domains (Extended Data Fig. 7c).  
 120 Conversely, both *PEAR1* and *TMO6* transcription were highly reduced in the procambial tissue of  
 121 cytokinin signalling loss-of-function mutant *wooden-leg (wol)*<sup>5,19</sup> (Fig. 3b and Extended Data Fig. 7d)  
 122 and in plants overexpressing *ARR22*<sup>20</sup>, an inhibitor of cytokinin signalling (Fig. 3d and e). However,  
 123 expression of both genes was restored by the induction of cytokinin signalling in *wol* (Fig. 3c and  
 124 Extended Data Fig. 7d). In addition, we validated the requirement of cytokinin signalling for *PEAR1*

expression during embryogenesis (Extended Data Fig. 7e-r, and see Supplementary Information). Taken together, our results indicate that initiation of *PEAR1* expression in early embryogenesis is independent of cytokinin signalling, but by the time the bisymmetric cytokinin pattern is formed at early heart stage, *PEAR1* transcription is activated and maintained post-embryonically by cytokinins.

Almost no periclinal cell divisions were observed in the cells non-adjacent to the pericycle, including PSE-IN where both cytokinin response and PEAR protein are present (Fig. 1b-d and 3a), suggesting an inhibitory mechanism restricts PEAR function in the inner cells. We previously observed an increased cell number in the vascular tissue of quadruple loss-of-function mutant of the five *Class III HOMEODOMAIN LEUCINE ZIPPER (HD-ZIP III)* genes<sup>21,22</sup> (Fig. 4a, d and g). These ectopic cell divisions occur in cells non-adjacent to the pericycle (Extended Data Fig. 8a-e). We observed high levels of three HD-ZIP III proteins, PHABULOSA (PHB), CORONA (CNA) and REVOLUTA (REV), in non-dividing procambial cells, IPC and PSE-IN, whereas their expression was absent in the actively dividing cells of the PSE and PSE-LN (Extended Data Fig. 8f-k). In this domain endodermal-derived mobile miR165/6 eliminates *HD-ZIP III* messenger RNA<sup>22,23</sup>, suggesting that HD-ZIP III inhibit periclinal cell divisions of PSE-IN by antagonizing the functions of co-localized PEAR proteins. This is further supported by our observation that overexpression of *PEAR1* in the miRNA-resistant *phb-1d* mutant which has elevated levels of PHB<sup>22,23</sup> is less effective than overexpressing *PEAR1* in wild-type plants (Extended Data Fig. 8l-o). Hence, to sharpen the boundary between dividing and non-dividing cells, the expression pattern of both HD-ZIP III and PEAR proteins must be tightly controlled.

Auxin is known to promote the xylem associated *HD-ZIP III* transcription<sup>24,25</sup>. However, *PHB*, *CNA* and *REV* show broader expression of both transcriptional and translational reporters (Fig. 4h, Extended Data Fig. 8f-h and Extended Data Fig. 9), suggesting that other factors may enhance *HD-ZIP III* transcription in the peripheral region. Interestingly, we observed a significant reduction of *CNA* transcription in PSE-neighbouring cells in the *pear* quintuple background (Fig. 4h-j). In addition, *PEAR1* overexpression enhanced the transcription of *HD-ZIP III* genes, especially in the central domain of vascular tissue (Extended Data Fig. 9). These data suggest that PEAR1 locally enhances *HD-ZIP III* transcription at PSE-neighbouring cells. As previous work has reported that PEAR1 has the potential to bind *HD-ZIP III* promoters<sup>26,27</sup>, it is possible that these interactions are direct. As *HD-ZIP III* and *PEAR1* show complementary expression patterns, we explored whether HD-ZIP III could regulate *PEAR1* transcription. *PEAR1* expression was severely attenuated in mutants showing elevated levels of HD-ZIP III such as *phb-1d* and *shr-222* (Extended Data Fig. 8p-t). Together these data suggest a feedback loop between *HD-ZIP III* and *PEAR1* transcription.

Furthermore, to examine a possible effect of the HD-ZIP III on the mobile PEAR1 proteins, we measured the diffusion coefficient and movement pattern of PEAR1-GFP in wild type and in the *hd-zip III* quadruple mutant where PSE is formed in a triarch arrangement but *PEAR1* transcription is restricted to PSE as observed in wild type (Extended Data Fig. 8u and w). We observed that the diffusion coefficient of PEAR1-GFP is significantly higher and the protein moves further in the mutant compared to wild type (Fig. 4k-m and Extended Data Fig. 4). To understand the significance of this enhanced PEAR1 movement, we analysed the cell proliferation pattern of combinatorial *pear1 pear2 dof6 obp2 hca2 phb phv cna athb8* nonuple and *pear1 pear2 dof6 tmo6 obp2 hca2 phb phv cna athb8* decuple loss-of-function mutants. We found that these mutants showed a reduced number of periclinal cell divisions in the vascular cells both adjacent to and non-adjacent to the pericycle compared to the

*hd-zip III* quadruple mutant (Fig. 4a-g and Extended Data Fig. 8d-e). This indicates that HD-ZIP III inhibit periclinal cell division partially through inhibiting PEAR1 movement to position the cell division zone around phloem.

In order to further conceptualize the observed interactions between PEAR and HD ZIP III and test the capacity of this network to generate sharp boundaries, we incorporated the PEAR factors into a spatially one dimensional network model with HD-ZIP III, miR165/6, auxin and cytokinin as defined in previous theoretical studies<sup>8,28,29</sup> (Supplementary Modelling Information). The model is defined on a line in one spatial dimension representing 3, 4 or 5 cells from the centre of the xylem axis to the outer edge of the PSE cell (Extended Data Fig. 10a-d and Supplementary Modelling Information). One particularly interesting aspect of the system is that the network involves dual negative feedback loops, in which *HD-ZIP III* transcription is activated by PEAR1 (Interaction (1) in Extended Data Fig.10c), while in turn both *PEAR1* transcription and protein movement are inhibited by HD-ZIP III (Interaction (2) and (3), respectively in Extended Data Fig.10c). We ran simulations exploring the steady state patterns created in networks with the above interactions and in scenarios when one of the interactions was missing (Extended Data Fig. 10d-h). Based solely on two inputs imposed at the margins, auxin (xylem) and miR165/6 (outer margin), the model predicts the spatial distribution of cytokinin, as well as PEAR and HD-ZIP III proteins (Extended Data Fig. 10d-h). The version of the model incorporating all three interactions (i.e repressing both the transcription and movement of PEAR) results in the sharpest gradients of PEAR and HD-ZIP III proteins (Extended Data Fig. 10f) with both PEAR1 protein and HD-ZIP III localized within the PSE-IN, consistent with experimental observations (Extended Data Fig. 10e). To our knowledge this is the first report of a role for the dual regulation of both transcription and movement of a developmental regulator in sharpening boundaries.

Collectively our research has uncovered a regulatory network involving the dual regulation of gene transcription and protein movement, in which the spatial distribution of phytohormones and small RNA is decoded into the activity of two functionally antagonistic sets of transcription factors, PEAR and HD-ZIP III, during root procambial development (Extended Data Fig. 10i). The mobile PEAR factors promote cell proliferation around the two early protophloem sieve element cell files, which constitute two new organizers just proximal to the quiescent centre. These organizers surround a more quiescent central zone defined by the HD-ZIP III factors. In this way, the PEAR - HD-ZIP III module specifies a lateral meristem within an apical meristem and as such, forms a foundation for further cambial development<sup>30</sup>. Therefore, in the future it will be interesting to determine how extensively this procambial pathway also contributes to ontogenetically late processes such as wood and storage organ formation in the crop species.

## Acknowledgements

We thank E. Scarpella, ABRC and NASC for materials, Katja Kainulainen, Karolina Blajecka, Mikko Herpola and Alessandro Mainardi for technical assistance, Natalie Clark for assisting with the scanning FCS analysis, Jenny Jansen for technical assistance with microarray hybridizations, Oleg Kambur and Lothar Kalmbach for assistance in generating the heatmap, and D. Weijers and T. Kakimoto for helpful discussions, A. Groenheide, E. Cornelissen, M. Chu, A. Korppoo and Alba Rodriguez Diez for technical support. This work was supported by Finnish Centre of Excellence in Molecular Biology of

209 Primary Producers (Academy of Finland CoE program 2014-2019) decision #271832, the Gatsby  
 210 Foundation [GAT3395/PR3], University of Helsinki [award 799992091] and the European Research  
 211 Council Advanced Investigator Grant SYMDEV [No. 323052] to Y. H., a NSF-BBSRC MCSB  
 212 1517058 to R.S. and Y.H., an ERC Consolidator grant (PLANTSTEMS), a Heisenberg Professorship  
 213 of the German Research Foundation (DFG, GR2104/5-1) and the SFB 873 (DFG) to T.G., the  
 214 Netherlands Organization for Scientific Research (NWO; VIDI-864.13.001) and The Research  
 215 Foundation - Flanders (FWO; Odysseus II G0D0515N and 12D1815N) to W.S. and B.D.R.,  
 216 respectively, a JSPS postdoctoral fellowship for research abroad and JSPS KAKENHI Grant  
 217 [17K15138] to S.M., Swiss National Science Foundation Postdoc Mobility Grant (P300P3\_147894)  
 218 to P.R., a JSPS Research Fellowship for Young Scientists and JSPS KAKENHI Grant [JP16J00131] to  
 219 K.T., Bayer Science and Education Foundation, German Academic Exchange Service (DAAD) to B.B.,  
 220 The Finnish Academy of Science to J. O. H., Herchel Smith Postdoctoral Research Fellowship  
 221 (Herchel Smith Fund) to S.O.

222

223

## 224 **Author contributions**

225 S.M., P.R. and I.S. contributed equally to this work. S.M. characterized the molecular interactions  
 226 among PEAR and HD-ZIP III module. P.R. identified and quantified phenotype in the PEAR loss of  
 227 function mutants with help of B.B. I.S. determined phloem specific DOFs and their downstream genes  
 228 with input from B.D.R., W.S., M.B. and G. H. K.T. characterized *PEAR-HD-ZIP III* combinatorial  
 229 mutants. B.B. generated *tmo6* CRISPR mutants. J.H. performed in situ hybridization. N.M. and A.B.  
 230 designed and performed computational modelling. H.H. produced CRE1 inducible line. S.O. assisted  
 231 in the microarray experiments. K.H. and K.N. produced HD-ZIP III reporter lines. O.S. and A.P.M.  
 232 provided the destination vector pSm43GW. R.S. and A.P.M. provided the *pARR5::RFP* line. E.S.W.,  
 233 Y.K., T.G. and C.M. shared informative non-published data. R.S. analysed diffusion coefficient of  
 234 PEAR1-GFP with P.R. B.D.R. and Y.H. participated in experimental design. S.M. and Y.H. wrote the  
 235 manuscript and all authors commented on the manuscript. B.D.R. and Y.H. are co-corresponding  
 236 authors.

237

238

## 239 **References**

240

- 241 1 Esau, K. *Vascular differentiation in plants*. (Holt, Rinehart and Winston, 1965).
- 242 2 Crick, F. H. & Lawrence, P. A. Compartments and polyclones in insect development. *Science* **189**,  
 243 340-347 (1975).
- 244 3 Dickson, B., Sprenger, F. & Hafen, E. Prepattern in the developing Drosophila eye revealed by an  
 245 activated torso--sevenless chimeric receptor. *Genes Dev* **6**, 2327-2339 (1992).
- 246 4 McConnell, J. R. *et al.* Role of PHABULOSA and PHAVOLUTA in determining radial patterning  
 247 in shoots. *Nature* **411**, 709-713, doi:10.1038/35079635 (2001).
- 248 5 Mähönen, A. P. *et al.* A novel two-component hybrid molecule regulates vascular morphogenesis of  
 249 the *Arabidopsis* root. *Genes Dev* **14**, 2938-2943 (2000).
- 250 6 Mähönen, A. P. *et al.* Cytokinin signaling and its inhibitor AHP6 regulate cell fate during vascular

development. *Science* **311**, 94-98, doi:10.1126/science.1118875 (2006).

7 Bishopp, A. *et al.* A mutually inhibitory interaction between auxin and cytokinin specifies vascular pattern in roots. *Curr Biol* **21**, 917-926, doi:10.1016/j.cub.2011.04.017 (2011).

8 De Rybel, B. *et al.* Plant development. Integration of growth and patterning during vascular tissue formation in *Arabidopsis*. *Science* **345**, 1255-1259, doi:10.1126/science.1255215 (2014).

9 Vatén, A. *et al.* Callose biosynthesis regulates symplastic trafficking during root development. *Dev Cell* **21**, 1144-1155, doi:10.1016/j.devcel.2011.10.006 (2011).

10 Brady, S. M. *et al.* A high-resolution root spatiotemporal map reveals dominant expression patterns. *Science* **318**, 801-806, doi:10.1126/science.1146265 (2007).

11 Yanagisawa, S. The Dof family of plant transcription factors. *Trends Plant Sci* **7**, 555-560 (2002).

12 Kim, H. S. *et al.* The DOF transcription factor Dof5.1 influences leaf axial patterning by promoting *Revoluta* transcription in *Arabidopsis*. *Plant J* **64**, 524-535, doi:10.1111/j.1365-313X.2010.04346.x (2010).

13 Skirycz, A. *et al.* DOF transcription factor AtDof1.1 (OBP2) is part of a regulatory network controlling glucosinolate biosynthesis in *Arabidopsis*. *Plant J* **47**, 10-24, doi:10.1111/j.1365-313X.2006.02767.x (2006).

14 Rueda-Romero, P., Barrero-Sicilia, C., Gómez-Cadenas, A., Carbonero, P. & Oñate-Sánchez, L. *Arabidopsis thaliana* DOF6 negatively affects germination in non-after-ripened seeds and interacts with TCP14. *J Exp Bot* **63**, 1937-1949, doi:10.1093/jxb/err388 (2012).

15 Guo, Y., Qin, G., Gu, H. & Qu, L. J. Dof5.6/HCA2, a Dof transcription factor gene, regulates interfascicular cambium formation and vascular tissue development in *Arabidopsis*. *Plant Cell* **21**, 3518-3534, doi:10.1105/tpc.108.064139 (2009).

16 Schlereth, A. *et al.* MONOPTEROS controls embryonic root initiation by regulating a mobile transcription factor. *Nature* **464**, 913-916, doi:10.1038/nature08836 (2010).

17 Wallner, E. S. *et al.* Strigolactone- and Karrikin-Independent SMXL Proteins Are Central Regulators of Phloem Formation. *Curr Biol* **27**, 1241-1247, doi:10.1016/j.cub.2017.03.014 (2017).

18 Siligato, R. *et al.* MultiSite Gateway-Compatible Cell Type-Specific Gene-Inducible System for Plants. *Plant Physiol* **170**, 627-641, doi:10.1104/pp.15.01246 (2016).

19 Mähönen, A. P. *et al.* Cytokinins regulate a bidirectional phosphorelay network in *Arabidopsis*. *Curr Biol* **16**, 1116-1122, doi:10.1016/j.cub.2006.04.030 (2006).

20 Kiba, T., Aoki, K., Sakakibara, H. & Mizuno, T. *Arabidopsis* response regulator, ARR22, ectopic expression of which results in phenotypes similar to the *wol* cytokinin-receptor mutant. *Plant Cell Physiol* **45**, 1063-1077, doi:10.1093/pcp/pch128 (2004).

21 Prigge, M. J. *et al.* Class III homeodomain-leucine zipper gene family members have overlapping, antagonistic, and distinct roles in *Arabidopsis* development. *Plant Cell* **17**, 61-76, doi:10.1105/tpc.104.026161 (2005).

22 Carlsbecker, A. *et al.* Cell signalling by microRNA165/6 directs gene dose-dependent root cell fate. *Nature* **465**, 316-321, doi:10.1038/nature08977 (2010).

23 Miyashima, S., Koi, S., Hashimoto, T. & Nakajima, K. Non-cell-autonomous microRNA165 acts in a dose-dependent manner to regulate multiple differentiation status in the *Arabidopsis* root. *Development* **138**, 2303-2313, doi:10.1242/dev.060491 (2011).

24 Baima, S. *et al.* The expression of the *Athb-8* homeobox gene is restricted to provascular cells in

- 293 *Arabidopsis thaliana*. *Development* **121**, 4171-4182 (1995).
- 294 25 Donner, T. J., Sherr, I. & Scarpella, E. Regulation of preprocambial cell state acquisition by auxin  
295 signaling in *Arabidopsis* leaves. *Development* **136**, 3235-3246, doi:10.1242/dev.037028 (2009).
- 296 26 O'Malley, R. C. *et al.* Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape.  
297 *Cell* **166**, 1598, doi:10.1016/j.cell.2016.08.063 (2016).
- 298 27 Gaudinier, A. *et al.* Enhanced Y1H assays for *Arabidopsis*. *Nat Methods* **8**, 1053-1055,  
299 doi:10.1038/nmeth.1750 (2011).
- 300 28 Muraro, D. *et al.* Integration of hormonal signaling networks and mobile microRNAs is required  
301 for vascular patterning in *Arabidopsis* roots. *Proc Natl Acad Sci U S A* **111**, 857-862,  
302 doi:10.1073/pnas.1221766111 (2014).
- 303 29 Mellor, N. *et al.* Theoretical approaches to understanding root vascular patterning: a consensus  
304 between recent models. *J Exp Bot* **68**, 5-16, doi:10.1093/jxb/erw410 (2017).
- 305 30 De Rybel, B., Mähönen, A. P., Helariutta, Y. & Weijers, D. Plant vascular development: from early  
306 specification to differentiation. *Nat Rev Mol Cell Biol* **17**, 30-40, doi:10.1038/nrm.2015.6 (2016).

307  
308

## 309 Figure legends

310

### 311 Figure 1| Periclinal cell divisions are centred around PSE, a domain highlighted by mobile PEAR 312 transcription factors.

313 **a**, Schematic representation of procambial cells based on the position relative to PSE (red) and outer  
314 pericycle (gray). PSE neighbouring cells are classified as PSE lateral neighbour (PSE-LN, orange), a  
315 cell adjacent to both PSE and pericycle, or PSE internal neighbour (PSE-IN, dark-green), a cell  
316 adjacent to PSE but not pericycle. Intervening procambial cells are classified as outer PC (OPC,  
317 yellow), a procambial cell adjacent to pericycle, or internal PC (IPC), a procambial cell non-adjacent  
318 to pericycle. PX and MX represent protoxylem and metaxylem, respectively. **b**, Number of periclinal  
319 cell divisions in each cell during procambial development (273 division events in total from 13  
320 independent roots, also see Supplementary Information). PSE and PSE-LN exhibited higher  
321 proliferative activity. Bar graphs represent mean. Error bars are s.d. Dots, individual data points. **c**,  
322 Expression of *pPEAR1::GFP-GUS* ( $n=17$ ) exhibits a highly PSE-specific expression pattern in the  
323 vascular tissue, though a residual level of GFP signal is observed in PSE-IN, most likely due to the  
324 retention of fluorescent protein after the division of PSE. **d**, Expression of the translational fusion of  
325 PEAR1 to GFP ( $n=15$ ). Fluorescent signal is observed not only in PSE but also in its neighbouring  
326 cells, including PSE-LN and PSE-IN. In **c** and **d**,  $n$  represents independent biological samples. White,  
327 dark-green, orange arrowheads and asterisks indicate PSE, PSE-IN, PSE-LN and protoxylem,  
328 respectively. Scale bars, 25  $\mu$ m.

329

### 330 Figure 2| *PEAR* genes activate periclinal cell division by controlling downstream genes in non- 331 cell autonomous manner.

332 **a-e**, Cross-section of wild type (**a**), *pear* quintuple (*pear1 pear2 dof6 obp2 hca2*) (**b**), *pear* triple (*pear1*  
333 *pear2 tmo6*) (**c**), *pear* quadruple (*pear1 pear2 dof6 tmo6*) (**d**) and *pear* hextuple (*pear1 pear2 dof6*  
334 *obp2 hca2 tmo6*) (**e**), respectively. Each image is representative of independent biological samples



analysed in **f. f**, Number of procambial and phloem cell files in wild type and *pear* combinatorial mutants. Values were calculated from root cross sections at the differentiation zone. Boxplot centres show median. For more information on boxplots, see Methods. Statistically significant differences between groups were tested using Tukey's HSD test  $p < 0.05$ . For individual *P* values, see Supplementary Table 3. *n*, independent biological samples. **g-h**, Expression of selected PEAR1/2 downstream genes in wild type, PEAR2 overexpression plant and *pear* hextuple mutant. *SMXL3* is expressed in phloem and procambial tissue, whose expression is induced by PEAR2 overexpression, but not altered in *pear* hextuple. *AT4G00950* gene is expressed in PSE and its neighbouring cells, whose expression is induced by PEAR2 overexpression and reduced in *pear* hextuple mutant. Number in each panel indicates samples with similar results of the total independent biological samples analysed. White arrowheads, PSE. Asterisks, protoxylem. Scale bars, 25  $\mu$ m.

**Figure 3| Cytokinin signalling triggers *PEAR1* expression.**

**a**, Expression of *ARR5* and *PEAR1* overlaps at the initial stage (**a'**) and early proliferative phase (**a''**), *n*=15. **b-c**, Transcription of *PEAR1* in *wol* root, which is conditionally rescued by *CRE1* induction (Est, estradiol treated). *PEAR1* transcription is severely reduced in the condition with attenuated cytokinin response (**b**, *n*=6), and is restored after three days induction of *CRE1* (**c**, arrowheads, *n*=6). **d-e**, *PEAR1* transcription is down-regulated after 48 hours of *ARR22* induction. *n*=5 and 7, respectively. In **a-e**, *n* represents independent biological samples. White arrowheads, PSE. Scale bars, 25  $\mu$ m.

**Figure 4| Antagonistic function of PEAR1 and HD-ZIP III sharpens the boundary between dividing and non-dividing cells.**

**a-f**, An optical cross-section image of vascular tissue in wild type (**a**), *pear* quintuple (**b**), *pear* hextuple (**c**), *hd-zip III* quadruple (**d**), *pear hd-zip III* nonuple (**e**) and *pear hd-zip III* decuple mutant (**f**). Asterisks indicate pericycle cells. Each image is representative of independent biological samples analysed in **g**. **g** Quantification of vascular cell number. In the analysis of *pear* quintuple and *pear hd-zip III* nonuple, a population of slowly elongating roots is selected as described in Fig. 2f. **h-i**, Expression of *CNA* transcriptional reporter in the control (**h**, the heterozygous *pear* quintuple, *n*=4) and *pear* quintuple background (**i**, *n*=3). **j**, Fluorescent level in PSE-IN is significantly reduced in *pear* quintuple. *n* represents individual measurements across 4 or 3 independent biological samples, respectively. **k-l**, PEAR1-GFP localization in wild type (**k**, *n*=19) and *hd-zip III* quadruple mutant (**l**, *n*=17). PEAR1-GFP is broadly localized even in IPC in *hd-zip III* quadruple (**l**, light-green arrowheads). **m**, Average diffusion coefficient of PEAR1-GFP in wild-type and *hd-zip III* quadruple roots obtained by performing Raster Image Correlation Spectroscopy (RICS). In **g-i** and **k-m**, *n* represents independent biological samples. In **g**, boxplot centres show median. Statistically significant differences were tested using Tukey's HSD test  $p < 0.05$ . For individual *P* values, see Supplementary Table 3. In **j** and **m**, bar graphs represent mean. Error bars are s.d. (**j**) or s.e.m. (**m**). *P* values were calculated by two-sided Student's t-test (**j**) or Mann-Whitney *U* test (**m**). Dots, individual data points. White, dark-green, orange, light-green arrowheads indicate PSE, PSE-IN, PSE-LN and IPC, respectively. Scale bars, 25  $\mu$ m.

377

378 **Methods**

379

380 No statistical methods were used to predetermine sample size. The experiments were not randomized,  
 381 and investigators were not blinded to allocation during experiments and outcome assessment.  
 382 Experiments were repeated at least twice. All experiments were repeated successfully.

383

384 **Plant materials and growth condition**

385 *Arabidopsis thaliana* lines used in this study were either in Columbia or Landsberg erecta background.  
 386 The following alleles were obtained from the publicly available collections: *pear1* (CSHL\_GT8483)  
 387 in Ler, *pear2* (SALK\_088165) in Col-0, *obp2* (SK24984) in Col-4, *dof6* (Wiscseq\_Ds\_Llox351c08)  
 388 in Col-0, *hca2* (GK-466B10) in Col-0. Knock-out alleles of *TMO6* were generated using CRISPR-  
 389 Cas9 technology as previously described<sup>31</sup>. The following protospacer target sequence was selected as  
 390 it had no predicted off-site targets and allowed screening via *NheI* restriction using the CRISPR-P web  
 391 tool<sup>32</sup>. The Protospacer adjacent motif is underlined: GGACACCTGAGAGCTAGCTCCGG.  
 392 Successful mutagenesis was confirmed via Sanger sequencing in plants of the T2 and T3 generation  
 393 that no longer carried the Cas9 transgene. Four *TMO6* mutant alleles were identified: *tmo6-1* (+A),  
 394 *tmo6-2* (+T), *tmo6-3* (deletion of 5 bp and at the same time insertion of 26 bp) and *tmo6-4* (-5 bp)  
 395 (Extended Data Fig. 5a). The alleles *tmo6-1*, -2, and -3 were found in the *pear* hextuple mutant and  
 396 caused the *pear* hextuple phenotype, while *tmo6-4* was found in the *tmo6* single mutant, respectively.  
 397 The genotyping primers for these mutants are listed in Supplementary Table 1. *hd-zip III* quadruple  
 398 (*phb phv cna athb8*) was described previously<sup>21</sup>. Plant growth conditions were described previously<sup>5</sup>.

399

400 **Histological analysis**

401 Primary roots of vertically grown 4 to 5-day-old seedlings were used for histological analyses. For  
 402 confocal imaging, root samples were stained with propidium iodide (PI), aniline blue (AB) or SCRI  
 403 Renaissance 2200 (SR2200) (Renaissance Chemicals, UK). The method of PI and AB staining were  
 404 described previously<sup>9,33</sup>. For SR2200 stain, root samples were fixed in SR2200 solution (4%  
 405 paraformaldehyde, 0.1% (v/v) SR2200 in PBS buffer (pH7.4)). Then samples were washed with PBS  
 406 buffer and transferred into the ClearSee solution<sup>34</sup>. Confocal imaging was performed on Leica TCS  
 407 SP5, Leica TCS SP8, Leica TCS SP5 II HCS-A or Nikon C2 CLSM using a solid state blue laser  
 408 (480nm) for GFP, a green laser (514nm) for VENUS, a lime laser (DPSS 561nm) for RFP and PI, and  
 409 a UV laser (diode 405nm) for SR2200. Transverse plastic sections of root were performed as described  
 410 previously<sup>5</sup>. For histological analyses of embryo, dissected embryos were mounted in SR2200 solution  
 411 and visualized by the confocal microscopy.

412

413 **Mapping of the position of periclinal cell divisions**

414 A series of 2D confocal images of *Arabidopsis* root vascular tissue were recorded at 0.5  $\mu$ m intervals  
 415 using Nikon C2 CLSM or Leica TCS SP8. Cross section images in each developmental stage were  
 416 created by ImageJ software from a series of 2D confocal images, and the cell segmentation was done  
 417 using CellSeT<sup>35</sup>. For more information, see Supplementary Information.

418

## Box plots

Box plots were created with standard box blot setting (the first and third quartiles, split by the median; whiskers extend to a maximum of 1.5× interquartile range (IQR) beyond the box.) Outliers are indicated as black dots.

## DNA constructs and transgenic plants

Most of transgenic constructs were produced by using Gateway or multisite Gateway system (Invitrogen) as described previously<sup>18</sup>. To generate the transcriptional fusion constructs with GFP-GUS each promoter sequence was cloned into pDONR221 and fused to GFP-GUS coding sequence in the destination vector pBGWFS7 by normal LR reaction. For other transcriptional fusion constructs, including *pPEAR1::VENUSer*, *pPEAR2::VENUSer*, *pAHA3::RFPper*, *pOBP2::VENUSer*, *pHCA2::RFPper*, *pTMO6::RFPper* and *pREV::RFPper* and the transcriptional fusion constructs of PEAR1/PEAR2 downstream genes, each promoter was cloned into pDONRP4\_P1R, and assembled with the coding sequence of fluorescent reporter (VENUSer or RFPper) and terminator into the destination vectors, pHm43GW (Hygromycin resistant), pBm43GW (Basta resistant) or by multisite Gateway system. To produce the transcriptional fusion constructs of HD-ZIP III, including PHB and CNA, each promoter was inserted upstream of the GAL4:VP16 (GV) coding region of pBIB-UAS-GFPper-NtADH5'-GV vector<sup>36</sup>. For most of the translational fusion constructs of PEAR genes, except for *pPEAR1::PEAR1-GFP*, each promoter was cloned into the first-box vector pDONRP4\_P1R, and each coding sequence was cloned into vector pDONR221, thereafter each promoter and coding sequence were assembled with pDONR P2R\_P3-terminator/reporter into pHm43GW, pBm43GW or pFR7mGW by multisite Gateway system<sup>18</sup>. To generate other translational fusion constructs, including *pPEAR1::PEAR1-GFP*, *pCNA::CNA-GFP*, *pATHB8::ATHB-GFP* and *pREV::REV-GFP*, each genomic fragment which contains promoter, coding and its 3' region, was cloned into pAN19 vector. Then GFP coding sequence was fused to C-terminus of each coding sequence. Finally, each translational fusion sequence was inserted into the modified pBIN19 vector with Basta resistance<sup>23</sup>. For the overexpression construct, including *PEAR* genes and *CRE1*, the coding sequence of each genes was assembled with stele-specific estradiol-inducible promoter (pCRE1[XVE]) into pHm43GW or pBm43GW by the Multisite Gateway system described previously<sup>18</sup>. To construct *pPEAR1[XVE]::icals3m*, 1.5kb *PEAR1* promoter was cloned into p1R4-ML:XVE vector<sup>18</sup>, and assembled with *icals3m* sequence into pBm43GW<sup>9</sup>. The primers for DNA construction and the list of plasmids are shown in Supplementary Table 1.

## In situ hybridization

Amplified fragments of *PEAR1*, *PEAR2* and *OBP2* were cloned into pGEM-T Easy (Promega) vector and fragments of *DOF6*, *HCA2*, *TMO6* into pCR-Bunt II-TOPO vector (Invitrogen) following manufacturer's instructions. In order to obtain antisense probes, plasmids were first linearized by restriction enzyme treatment: *MluI* for *PEAR1* and *OBP2*, *ScaI* for *PEAR2*, *HindIII* for *TMO6* and *DOF1*, and *XbaI* for *HCA2* were used. Linearized plasmids were digoxigenin (DIG) labelled using DIG RNA Labelling Kit (Roche) following manufacturer's instructions. For *PEAR1*, *OBP2*, *TMO6* and *DOF1*, T7 RNA polymerase and for *PEAR2* and *HCA2*, SP6 RNA polymerase were used. mRNA detection on a whole-mount seedlings was performed as described<sup>37</sup>. Images were taken with Zeiss

461 Axioimager microscope with either 20x or 40x objective.

462

### 463 **Transcriptome analysis**

464 Targets of PEAR1 and PEAR2 were identified by analysing transcriptional changes after  
465 dexamethasone (DEX) treatment of *pRPS5A::PEAR1-GR* and *pRPS5A::PEAR2-GR*. To identify  
466 putative direct targets, DEX treatment was also performed with cycloheximide (CHX), which inhibits  
467 protein synthesis and therefore activation of indirect targets. 3-day-old seedlings were grown on  
468 control medium and transferred to medium containing 10  $\mu$ M DEX or 10  $\mu$ M DEX and 10  $\mu$ M CHX  
469 for 2h, after which root tips were collected and RNA extraction was performed. Total RNA (100 ng)  
470 was labelled using GeneChip WT PLUS Reagent Kit (Thermo Fisher Scientific) and hybridized to  
471 GeneChip Arabidopsis Gene 1.1 ST array plates (Affymetrix). Sample labelling, hybridization to chips,  
472 and image scanning were performed according to the manufacturer's instructions. Microarray analysis  
473 was performed as previously described to yield significantly up-regulated genes ( $>1.0$ -fold;  $P < 0.05$ )<sup>8</sup>.  
474 Venn diagram of significantly up-regulated genes was made using Venny 2.1 on-line program  
475 ([http://bioinfogp.cnb.csic.es/tools/venny\\_old/venny.php](http://bioinfogp.cnb.csic.es/tools/venny_old/venny.php)). Previously published root spatiotemporal  
476 expression data was used to make a heatmap to visualize predicted expression patterns of all PEAR1  
477 and PEAR2 targets<sup>10</sup>. To have relative expression values for every gene in different root cell types and  
478 developmental stages, values for every gene were normalized based on its highest expression in one  
479 of the cell types. Heatmap was generated using R with gplots R-package<sup>38</sup>. The transcriptomics data  
480 files are submitted to GEO (accession number GSE115183).

481

### 482 **Reporter analysis of PEAR1/2 downstream genes**

483 When selecting genes for reporter analysis, putative direct targets were preferred. Significantly more  
484 direct targets were identified for *PEAR2*, and therefore those are overrepresented. Other considerations  
485 were how strongly they were upregulated, as well as their predicted expression pattern. Expression in  
486 early procambium or early phloem and procambium was preferred. *AT1G49230*, *AT1G15080*,  
487 *AT3G16330*, *AT4G00950* and *SMXL3* are putative direct targets of PEAR2 and with predicted  
488 expression in early phloem/procambium. *AT3G54780* was chosen because it is a putative direct target  
489 of both PEAR1 and PEAR2, although no predicted expression data was available. *AT1G09460*, a direct  
490 target of PEAR2 and a target of PEAR1, was chosen because it was induced very strongly by both  
491 genes, although predicted to be expressed only very weakly in phloem/procambium.

492

### 493 **Quantitative RT-PCR analysis**

494 qRT-PCR analyses were performed as described previously<sup>39</sup>. Cytokinin treatment was done with  
495 10 $\mu$ M 6-Benzylaminopurine (BA), and experiments were performed in three biological repeats and  
496 each of these with 3 technical repeats. RNA was extracted with the RNeasy kit (QIAGEN). Poly(dT)  
497 cDNA was prepared from 1  $\mu$ g of total RNA with an iScript cDNA Synthesis Kit (Biorad) and analysed  
498 on a CFX384 Real-Time PCR detection system (BioRad) with iQ SYBR Green Supermix (BioRad)  
499 according to the manufacturer's instructions. Expression levels were normalized to those of *EEF1 $\alpha$*   
500 and *CDKA1;1*. The primers are listed in Supplementary Table 1.

501

### 502 **Phloem transport assay**

The phloem translocation was judged by the transport and unloading of 5(6) Carboxyfluorescein diacetate (CFDA) as describe<sup>40</sup>. After application of the dye, plants were kept in agar plates and only placed on regular cover slips for imaging.

### **Raster image correlation spectroscopy (RICS)**

To determine the rate of movement of GFP-labeled PEAR1 protein in wild type and *hd-zip III* quadruple (*phb phv cna athb8*) mutant background, Raster image correlation spectroscopy (RICS) was performed according to previous work<sup>41-43</sup>. Images were collected using a Zeiss 880 confocal microscope. Frames of 256x256 pixels were acquired using a raster scan with a dwell time of 8.19  $\mu$ sec pixel<sup>-1</sup> at a pixel size of either 100nm for 100 frames resulting in a line scan of 5.035ms. Diffusion coefficients were derived using the SimFCS software (<https://www.lfd.uci.edu/globals/>)<sup>44</sup> from GFP-labeled PEAR1 vascular cells within the first 70  $\mu$ m from the QC. Specifically, 18 observations from WT and 30 for the *hd-zip III* quadruple (*phb phv cna athb8*) mutant background were used for the RICS analysis using the SimFCS software. The RICS algorithm by comparing the intensity fluctuations of one pixel to the fluctuations of the pixels next to it and the fluctuations of one pixel to itself over time, produces a spatio-temporal Auto Correlation Function (ACF) that captures the fluorescence dynamics of the particles in the volume<sup>44,45</sup>. The ACF is decomposed into two correlation functions that depend on  $\xi$  (the spatial lag in  $x$ ) and  $\psi$  (the spatial lag in  $y$ ). The first correlation function,  $S(\xi, \psi)$ , calculates the spatio-temporal correlation due to the scanning of the microscope. The second correlation function,  $G(\xi, \psi)$ , calculates the spatio-temporal correlation due to particles diffusing in the medium. The ACF,  $G_S(\xi, \psi)$ , takes both of these correlations into account by multiplying them:  $G_S(\xi, \psi) = S(\xi, \psi) * G(\xi, \psi)$ . The functions are constructed assuming that the distribution of fluorescence intensities follows a 3D Gaussian distribution. The decomposition of the ACF into two parts allows RICS to distinguish random, Brownian motion from diffusing particles in the medium<sup>45</sup>. The software fits the RICS-ACF using the pixel dwell time, pixel size, line scan and the Point Spread Function (PSF) bean waist of 0.241nm as previously obtained<sup>41</sup> and returns the diffusion coefficient of the protein. The diffusion coefficient returned results in the ACF curve that best fits the data. Goodness of fit is determined by comparing the residuals to the amplitude of the ACF<sup>41-45</sup>.

### **Mathematical model**

The mathematical model is formulated as a set of ordinary differential equations describing the set of interactions shown in Figure 4o, defined on a one-dimensional array of discrete spatial compartments representing a cross-section of root tissue. The spatial subdivisions may represent either cell or cell wall compartments, with multiple compartments per cell so that intracellular resolution is present within the model. Three, four or five cells are simulated, from the centre of the stele at the xylem axis to the edge of the stele where phloem is formed. The model is implemented as a single stand-alone text file using Python 2.7 plus the open source libraries Scipy, from which the 'odeint' function was used to solve the differential equations, and Matplotlib, which was used to plot the figures. See Supplementary Modelling Information for more details.

### **Code availability**

545 The code for mathematical model is available on request.

546

547

#### 548 **Data availability**

549 All lines and data supporting the findings of this study are available from the corresponding author  
550 upon request. The microarray data files are available at Gene Expression Omnibus (GEO) (accession  
551 no. GSE115183).

552

553

#### 554 **Methods references**

555

556 31 Fauser, F., Schiml, S. & Puchta, H. Both CRISPR/Cas-based nucleases and nickases can be used  
557 efficiently for genome engineering in *Arabidopsis thaliana*. *Plant J* **79**, 348-359,  
558 doi:10.1111/tpj.12554 (2014).

559 32 Lei, Y. *et al.* CRISPR-P: a web tool for synthetic single-guide RNA design of CRISPR-system in  
560 plants. *Mol Plant* **7**, 1494-1496, doi:10.1093/mp/ssu044 (2014).

561 33 Wysocka-Diller, J. W., Helariutta, Y., Fukaki, H., Malamy, J. E. & Benfey, P. N. Molecular analysis  
562 of SCARECROW function reveals a radial patterning mechanism common to root and shoot.  
563 *Development* **127**, 595-603 (2000).

564 34 Kurihara, D., Mizuta, Y., Sato, Y. & Higashiyama, T. ClearSee: a rapid optical clearing reagent for  
565 whole-plant fluorescence imaging. *Development* **142**, 4168-4179, doi:10.1242/dev.127613 (2015).

566 35 Pound, M. P., French, A. P., Wells, D. M., Bennett, M. J. & Pridmore, T. P. CellSeT: novel software  
567 to extract and analyze structured networks of plant cells from confocal images. *Plant Cell* **24**, 1353-  
568 1361, doi:10.1105/tpc.112.096289 (2012).

569 36 Hashimoto, K., Miyashima, S., Sato-Nara, K., Yamada, T. & Nakajima, K. Functionally Diversified  
570 Members of the MIR165/6 Gene Family Regulate Ovule Morphogenesis in *Arabidopsis thaliana*.  
571 *Plant Cell Physiol* **59**, 1017-1026, doi:10.1093/pcp/pcy042 (2018).

572 37 Hejatko, J. *et al.* In situ hybridization technique for mRNA detection in whole mount *Arabidopsis*  
573 samples. *Nat Protoc* **1**, 1939-1946, doi:10.1038/nprot.2006.333 (2006).

574 38 Wickham, H. & Sievert, C. *ggplot2 : elegant graphics for data analysis*. Second edition. edn,  
575 (Springer, 2016).

576 39 De Rybel, B. *et al.* A bHLH complex controls embryonic vascular tissue establishment and  
577 indeterminate growth in *Arabidopsis*. *Dev Cell* **24**, 426-437, doi:10.1016/j.devcel.2012.12.013 (2013).

578 40 Oparka, K. J., Duckett, C. M., Prior, D. A. M. & Fisher, D. B. Real - time imaging of phloem  
579 unloading in the root tip of *Arabidopsis*. *The Plant Journal* **6**, 759-766, doi:doi:10.1046/j.1365-  
580 313X.1994.6050759.x (1994).

581 41 Clark, N. M. *et al.* Tracking transcription factor mobility and interaction in *Arabidopsis* roots with  
582 fluorescence correlation spectroscopy. *Elife* **5**, doi:10.7554/eLife.14770 (2016).

583 42 Clark, N. M. & Sozzani, R. Measuring Protein Movement, Oligomerization State, and Protein-  
584 Protein Interaction in *Arabidopsis* Roots Using Scanning Fluorescence Correlation Spectroscopy  
585 (Scanning FCS). *Methods Mol Biol* **1610**, 251-266, doi:10.1007/978-1-4939-7003-2\_16 (2017).

586 43 O'Lexy, R. *et al.* Exposure to heavy metal stress triggers changes in plasmodesmatal permeability

via deposition and breakdown of callose. *J Exp Bot* **69**, 3715-3728, doi:10.1093/jxb/ery171 (2018).

44 Digman, M. A. *et al.* Measuring fast dynamics in solutions and cells with a laser scanning  
microscope. *Biophys J* **89**, 1317-1327, doi:10.1529/biophysj.105.062836 (2005).

45 Digman, M. A. & Gratton, E. Analysis of diffusion and binding in cells using the RICS approach.  
*Microsc Res Tech* **72**, 323-332, doi:10.1002/jemt.20655 (2009).

## Extended Data legends

### Extended Data Fig. 1| Quantification of periclinal cell division during procambial development.

**a**, Schematic representation of root vascular tissue of *Arabidopsis*. Procambial cells originate from their initial cells, and periclinal cell division increases the cell files during the proliferative phase, eventually resulting in a bisymmetric vascular pattern composed of a pair of phloem poles, which are separated from central xylem axis by intervening procambium. **b**, An example of mapping the position of periclinal cell divisions from the initial cells. From each position within the root vascular tissue (arrows), an optical cross-section image is constructed, and cells were segmented using CellSet. **c**, The number of periclinal cell divisions in each cell position (273 division events from 13 independent roots). **d**, The mean cell number in each category during procambial development. The number of events per cell in each group was calculated by dividing the number of events by the mean cell number of each group during development (See Supplementary Information).

### Extended Data Fig. 2| Inhibition of symplastic connection in early PSE results in the reduction of vascular cell number and in PSE-specific PEAR1-GFP localization.

**a**, Aniline blue stained primary root of *pPEAR1[XVE]::icals3m* after 24 hours of induction. Callose deposition occurs superficially in PSE cells (arrowheads,  $n=10$ ). **b-c**, The vascular tissue of *pPEAR1[XVE]::icals3m* root, in non-induced (**b**,  $n=13$ ) and after three days induction (**c**,  $n=9$ ). In non-induced condition, PSE cells (white arrowheads) and their neighbouring cells, composed of MSE (dark-green arrows) and two lateral companion cells (orange arrowheads), are spatially separated from the xylem axis by intervening procambium. By contrast, after three days induction of callose deposition in PSE cells, only a single SE cell file is formed in each phloem pole (**c**, white arrowheads), and its neighbouring cells often touch the xylem axis (**c**, yellow hashtags). Number of procambial and phloem cell files is significantly reduced after three days induction. Boxplot centres show median. For more information on boxplots, see Methods. *P* value was calculated by two-sided Student's t-test. **d-e**, Expression of *Sister of APL* (*SAPL*, *AT3G12730*) and *ATPase 3* (*AHA3*, *AT5G57350*) in *pPEAR1[XVE]::icals3m* before (**d**,  $n=10$  and 4, respectively) and after 24 hours of induction (**e**,  $n=10$  and 4, respectively). *SAPL* is expressed in CC and MSE in meristematic zone, and *AHA3* is expressed in differentiated CC (**d**). After induction, expression of these genes is restricted to a single cell file, indicating that symplastic cell communication between PSE and PSE-LN is required for the specification of PSE-neighbouring cell identity. **f-g**, PEAR1-GFP localization in *pCRE1[XVE]::icals3m* before (**f**,  $n=8$ ) and after 24 hours of induction (**g**,  $n=7$ ). PEAR1-GFP becomes specific to the PSE cell after the induction of callose deposition in whole vascular tissue, suggesting that PEAR1-GFP move in a short range via plasmodesmata. White, orange and dark-green arrowheads

629 indicate PSE, PSE-LN and PSE-IN, respectively. Asterisks indicate protoxylem (PX) cells. In **a-g**, *n*  
630 represents independent biological samples. Scale bars, 25  $\mu$ m.

631

632 **Extended Data Fig. 3| Identification of *PEAR* genes.**

633 **a**, *In silico* analysis of the early phloem abundant transcription factors. Nine transcription factors are  
634 shown to be expressed abundantly in the early phloem cell (S32 fraction), containing four types of  
635 transcription factors, including DOF-type, MADS-box, NAC-type and GATA-type transcription  
636 factors. **b**, A phylogenetic tree of 36 *Arabidopsis* DOF transcription factors is drawn using Clustal  
637 Omega (<https://www.ebi.ac.uk/Tools/msa/clustalo/>). **c**, Overexpression of *PEAR* genes, including  
638 *PEAR1*, *PEAR2*, *OBP2*, *DOF6*, *HCA2* and *TMO6*, under the *CRE1* inducible promoter enhances  
639 periclinal cell division in the vascular tissue. *n* represents independent biological samples. Bar graphs  
640 represent mean. Error bars are s.d. Dots, individual data points. *P* values were calculated by two-sided  
641 Student's t-test. **d**, Expression of six *PEAR* genes, including *PEAR1*, *PEAR2*, *DOF6*, *TMO6*, *OBP2*  
642 and *HCA2*, show similar expression patterns to *PEAR1*, in which both mRNA and transcriptional  
643 fusion reporter exhibit PSE-specific pattern with a broad protein localization. *HCA2* translational  
644 fusion in wild-type background exhibits weak but detectable signal in PSE-neighbouring cells (arrows),  
645 and its expression level is enhanced in the *pear* quintuple mutant background. Though *TMO6* mRNA  
646 is highly specific to PSE cells, its transcriptional fusion reporter shows a broad but PSE abundant  
647 expression pattern with a broad TMO6 protein localization. Mobility of TMO6 protein is more evident  
648 when TMO6-VENUS is expressed under PSE-specific *PEAR1* promoter (*pPEAR1::TMO6-VENUS*)  
649 in *pear* hextuple. Number in each panel indicates samples with similar results of the total independent  
650 biological samples analysed. Boxplot centres show median. *P* value calculated by two-sided Student's  
651 t-test. Dots, individual data points. **e**, A quantitative analysis of *PEAR* transcripts in *pear1 pear2* double  
652 mutant background. Note that the level of transcripts of three *PEAR* genes, including *TMO6*, *DOF6*  
653 and *HCA2*, is elevated in *pear1 pear2* background, suggesting that a compensation mechanism would  
654 mask the effect of *pear1 pear2* loss of function (also see Supplemental Information). Bar graphs  
655 represent mean. Error bars are s.d. Dots, individual data points. *P* values were calculated by two-sided  
656 Student's t-test. Scale bars, 25  $\mu$ m.

657

658 **Extended Data Fig. 4| PEAR1-GFP localization during procambial development.**

659 **a-d**, PEAR1-GFP localization in wild-type background (*n*=19, independent biological samples). The  
660 position of each optical section is indicated in the left panel showing the longitudinal section. At the  
661 position of the vascular initial cells, weak PEAR1-GFP signal is observed in PSE and neighbouring  
662 procambial cells but not in the xylem cells (**a**). During an early stage of the proliferative phase, the  
663 highest PEAR1-GFP signal is detected in the PSE, and substantial level of PEAR1-GFP signal is  
664 observed in PSE neighbouring cells, PSE-LN and PSE-IN (**b**, **c**) and its expression is maintained by  
665 the end of proliferation stage (**d**), indicating that the expression pattern of PEAR1-GFP is correlated  
666 with the domain having high proliferative activity, except for PSE-IN where almost no periclinal cell  
667 division is detected (see Fig. 1b). **e-h**, PEAR1-GFP localization in *hd-zip III* quadruple (*phb phv cna*  
668 *athb8*) mutant background (*n*=17, independent biological samples). The position of each optical  
669 section is indicated in left panel showing the longitudinal section. Broad localization of PEAR1-GFP  
670 is detected at the level of vascular initials. Central domain is highlighted with a dotted square (**e**). At



the early stage of proliferative stage, fluorescent signal is detected in IPC cells (light-green arrowheads), as well as PSE and its neighbouring cells (**f, g**), and gradually becomes specific to PSE and its neighbours (**h**). **i**, Quantification of PEAR1-GFP signal in each cell type. Fluorescent intensity of PEAR1-GFP in IPC and PSE-IN cells during proliferative phase (**b-c** in wild-type, **f-g** in *hd-zip III* quadruple, respectively) was measured and normalized to the fluorescent intensity in PSE cells, confirming a broad distribution of PEAR1-GFP in *hd-zip III* quadruple. *n* represents individual measurements across 5 (wild type) or 4 (*hd-zip III* quadruple) independent biological samples, respectively. Bar graphs represent mean. Error bars are s.d. Dots, individual data points. *P* values were calculated by two-sided Student's t-test. White, orange and dark-green and light-green arrowheads indicate PSE, PSE-LN, PSE-IN and IPC respectively. Scale bars represent 25  $\mu$ m.

#### **Extended Data Fig. 5| Loss of function of *PEAR* genes.**

**a**, Organization of *PEAR* genes and CRISPR/Cas9-induced mutation in TMO6 locus. Deletions are denoted by dashes; insertions and a replacement are indicated by red letters. **b**, Quantification of phloem and procambium cell files in lower (left) and higher (right) order *pear* combinatorial mutants. Tukey's HSD test is provided for all samples in Supplementary Table 3. **c**, *pear* quintuple mutant phenotype is suppressed by introduction of fluorescent-tagged PEAR proteins expressed under their native promoters. **d**, *pear* hextuple mutant phenotype is significantly suppressed by the introduction of PEAR1, DOF6 and TMO6 construct, but not PEAR2. In the *pear* hextuple background, *PEAR2* expression is highly reduced in the vascular tissue. **e**, Phloem unloading assay in wild type, *pear* quintuple (with shortest roots) and *pear* hextuple (*n*=15, 8 and 15, respectively). Fluorescent CFDA dye is loaded on the cotyledon and imaged two hours after application (see Methods). Phloem transport and unloading is not changed in the shortest roots of the *pear* quintuple mutant strongly affected in the radial growth. *pear* hextuple shows defects in phloem transport. **f**, Phenotype of *pear* hextuple mutant at the early developmental stage (1.5 days after germination). The cell number in vascular tissue of *pear* hextuple is significantly reduced before the onset of phloem PSE differentiation and activation of the phloem transport (see Supplementary Information). *P* value was calculated by two-sided Student's t-test. In **b-f**, *n* represents independent biological samples. In **b-d**, statistically significant differences between groups were tested using Tukey's HSD test *p*<0.05. For individual *P* values, see Supplementary Table 3. Boxplot centres show median. For more information on boxplots, see Methods. Scale bars represent 25  $\mu$ m.

#### **Extended Data Fig. 6| Identification of genes acting downstream of PEAR.**

**a**, Venn diagram showing the genes upregulated by overexpression of *PEAR1* or *PEAR2* with and without cycloheximide (chx). The analysis revealed 212 and 435 upregulated genes, in the respective experiments. Heatmap showing the predicted spatiotemporal expression patterns of all genes induced by PEAR1 or PEAR2. **b-i**, Expression patterns of eight selected genes responding to *PEAR2* overexpression. In control conditions, all genes exhibit a broad expression pattern, in which five of them are transcribed both in phloem and procambial cells (**b-d** and **i**), and the rest of them are in PSE and its surrounding cells where PEAR proteins are accumulated abundantly (**e-h**). Whereas expression of *SMXL3*, *ATIG09460* and *ATIG15080* are maintained even in *pear* hextuple (**b, c, f**), the expression level of five genes are attenuated (**d, e, g-i**). Number in each panel indicates samples with similar

713 results of the total independent biological samples analysed. **j**, Number of vascular cells after 3-days  
714 induction of overexpression of each *PEAR* downstream gene. In each case, several lines were analysed  
715 in parallel for a phenotypic change. Only *SMXL3* overexpression can increase the vascular cell number  
716 (confirmed in three independent lines). Boxplot centres show median. For more information on  
717 boxplots, see Methods. *n* represents independent biological samples. Dots, individual data points. *P*  
718 value was calculated by two-sided Student's t-test. Scale bars represent 25  $\mu$ m.

719

720 **Extended Data Fig. 7| Cytokinin controls *PEAR* expression.**

721 **a**, Expression of auxin (*pIAA2::GFP-GUS*) and cytokinin (*pARR5::RFP<sub>er</sub>*) response genes (*n*=15).  
722 Auxin response is restricted to the xylem cells at initial stage (**a'**) and maintained during development  
723 (**a''**, **a'''**). High cytokinin response is activated initially and maintained in PSE and its neighbouring  
724 cell (**a'**, **aa'**), and later becomes concentrated into the intervening procambial cells flanking xylem  
725 cells (**a'''**). **b**, Exogenous cytokinin application rapidly promotes the transcript level of *PEAR* genes,  
726 including *PEAR2*, *DOF6*, and *TMO6*. Asterisks indicate significant (*p*<0.05) upregulation as  
727 determined by a two-sided t-test on three biological replicates. Bar graphs represent mean. Error bars  
728 are s.e.m. For individual *P* values, see Supplementary Table 3. **c**, Sustained cytokinin application leads  
729 to ectopic transcription of *PEAR* genes. The optical cross section images are obtained after 4 days (for  
730 *PEAR1*, *PEAR2* and *HCA2*) or 1-day (*DOF6* and *OBP2*) treatment of 1 $\mu$ M of BA. **d**, Conditional  
731 induction of *CRE1* expression restores *TMO6* transcription in *wol* root. In the absence of cytokinin  
732 response, *TMO6* transcription is increased in pericycle and attenuated in the vascular tissue (Control,  
733 hashtags), and is restored in the vascular tissue after *CRE1* induction (Est). **e-l**, Expression pattern of  
734 auxin (*pIAA2::GFP-GUS*) and cytokinin (*pARR5::RFP<sub>er</sub>*) response reporters during embryogenesis  
735 in wild type (**e-h**, *n*=29, 13, 13, 11, respectively) and *wol* (**i-l**, *n*=12, 8, 6, respectively). At globular  
736 stage, auxin response is activated among provascular cells both in wild type (**e**) and in *wol* (**i**). At early  
737 heart stage, cytokinin response is activated in cells positioned below shoot apical meristem (**f**, ult,  
738 arrowheads), and the stripe of cytokinin response domain is formed by mid heart stage (**g**, ult,  
739 arrowheads), simultaneously auxin response becomes concentrated in the cells proximal to the  
740 cotyledon (**g**, ult, asterisks), resulting in the bisymmetric hormonal response pattern. During the  
741 torpedo stage, cytokinin response domain reaches to llt (**h**, llt, arrowheads). In *wol* embryos, activation  
742 of cytokinin response in vascular tissue does not occur and a radial auxin response pattern is maintained  
743 (**i-l**). **m-r**, Expression of *ARR5* and *PEAR1* during embryogenesis. In the wild-type embryo (**m-o**, *n*=17,  
744 25, 10, respectively), *PEAR1* is broadly transcribed among provascular cells both in ult and llt with  
745 radial symmetric pattern at the globular stage (**m**). At the heart stage, *PEAR1* transcription is enhanced  
746 in ult cells underneath the shoot apical meristem, which correlated with the activation of cytokinin  
747 response in this domain (**n**, arrowheads), and expression of both *ARR5* and *PEAR1* extends rootward  
748 and reaches to llt, becoming more concentrated within the cell files where phloem is specified post-  
749 embryonically (**o**, arrowheads). In *wol* embryos (**p-r**, *n*=19, 13, 13, respectively), *PEAR1* transcription  
750 is initiated among provascular cells at the globular embryo stage (**p**) similar to wild type (**m**), but  
751 neither cytokinin response nor *PEAR1* transcription occurs in ult at the heart stage (**q**), and *PEAR1*  
752 expression is gradually attenuated by the torpedo stage (**r**). ult and llt represent upper- and lower tier,  
753 respectively. In **a**, **e-r**, *n* represents independent biological samples. In **c** and **d**, number in each panel  
754 indicates samples with similar results of the total independent biological samples analysed. Scale bars

755 represent 25  $\mu$ m.

756

757 **Extended Data Fig. 8| HD-ZIP III restrict periclinal cell divisions during procambial**  
758 **development.**

759 **a-c**, Periclinal cell divisions in the cells non-adjacent to pericycle, including PSE-IN (**a**), IPC (**b**) and  
760 xylem cell (**c**), occurs in *hd-zip III* quadruple (*phb phv cna athb8*). *n*=8. **d-e**, The number of periclinal  
761 cell divisions in cells adjacent (**d**) or non-adjacent to pericycle (**e**) in wild type, *pear* and *hd-zip III*  
762 combinatorial mutants. In the analysis of *pear* quintuple and *pear hd-zip III* nonuple, a population of  
763 slowly elongating roots is selected as described in Fig. 2f. Boxplot centres show median. Statistically  
764 significant differences between groups were tested using Tukey's HSD test *p*<0.05. For individual *P*  
765 values, see Supplementary Table 3. **f-i**, Localization of PHB-GFP (**f**, *n*=12), CNA-GFP (**g**, *n*=10), REV-  
766 GFP (**h**, *n*=7) and ATHB8-GFP (**i**, *n*=5). **j-k**, Protein localization of PHB-GFP (**j**, *n*=7) and CNA-GFP  
767 (**k**, *n*=4) during procambial development. In the initial cells (**j'**, **k'**), both proteins are localized in  
768 metaxylem cells but not in PSE (white arrowheads). During the proliferation stage, PSE-IN (green  
769 arrowheads), which is produced by periclinal cell division in PSE, acquires the expression of both  
770 PHB-GFP (**j''**, **j'''**) and CNA (**k''**, **k'''**). **l-o**, Overexpression of PEAR1-VENUS under the *CRE1*  
771 inducible promoter in wild type (**l**, **n**, *n*=5, 3, respectively) and heterozygous *phb-1d* background (*phb-*  
772 *1d/+*, **m**, **o**, *n*=5, 5, respectively). After 18 hours of induction, PEAR1-VENUS signal is detected in  
773 both backgrounds (**l**, **m**), however, enhanced periclinal divisions are only observed in wild type (**l**),  
774 and not in *phb-1d* (**m**). Longer induction of *PEAR1* overexpression induces divisions even in *phb-1d*  
775 (**n**, **o**). **p-r**, *pPEAR1-PEAR1-GFP* expression is reduced in heterozygous *phb-1d/+* background. Most  
776 of *phb-1d* heterozygotes exhibit a single PEAR1-GFP expressing pole (**q**, 72% *n*=11), and the  
777 expression of PEAR1-GFP is almost completely abolished in some roots of *phb-1d* (**r**, 18% *n*=11). **s-**  
778 **t**, The expression of *pPEAR1::GFP* in wild type (**s**, *n*=10) and *shr-2* (**t**, *n*=9). The fluorescent signal  
779 is below the limit of detection in *shr-2*. **u-w**, Expression of *pPEAR1::GFP-GUS* in WT (**u**, *n*=19) and  
780 *hd-zip III* quadruple mutant (**w**, *n*=11). In **a-w**, *n* represents independent biological samples. White,  
781 orange and dark-green arrowheads indicate PSE, PSE-LN, PSE-IN respectively. Scale bars represent  
782 25  $\mu$ m.

783

784 **Extended Data Fig. 9| Overexpression of PEAR1 enhances the transcription of HD-ZIP III.**

785 The transcription patterns of four *HD-ZIP III*, including *PHB* (**a**, **b**), *CNA* (**c**, **d**), *REV* (**e**, **f**) and *ATHB8*  
786 (**g**, **h**) are visualized using their transcriptional fusion constructs. A longitudinal section is shown in the  
787 left panel, and the optical cross sections associated with this are shown in the right panels (the position  
788 of each section is indicated in the left panel). **a-b**, Transcription pattern of *PHB*  
789 (*pPHB::GV>UAS::GFP*) in *pCRE1[XVE]::PEAR1* plant before (**a**) and after 24 hours of induction  
790 of PEAR1 overexpression (**b**). *PHB* transcription is observed in whole vascular tissue at the initial and  
791 proliferative phase with peaks in xylem cells (**a'**), and its expression becomes concentrated into  
792 protoxylem cells (**a''** and **a'''**, asterisks indicate protoxylem cell). After the induction of PEAR1  
793 overexpression, *PHB* expression in the central domain of the vascular tissue is maintained at the later  
794 stage, resulting in the radially symmetric *PHB* transcription pattern (**b'** and **b''**). **c-d**, Transcription of  
795 *CNA* (*pCNA::GV>UAS::GFP*) in *pCRE1[XVE]::PEAR1-RFP* plant before (**c**) and after 24 hours of  
796 induction (**d**). *CNA* transcription is observed mainly in xylem lineage at initial cells (**c'**), and becomes

broader in whole vascular tissue, with peaks in procambial tissue, including PSE neighbouring cells (**c''**), and eventually its expression is gradually reduced in PSE and metaxylem, but is maintained in procambium, PSE neighbouring cells, as well as protoxylem cells (**c'''**). In a similar manner to *PHB*, *CNA* transcription in the central domain of the vascular tissue is maintained at the later stage when PEAR1-RFP is overexpressed (**d'''**). **e-f**, Transcription of *REV* (*pREV::RFP*) in *pCRE1[XVE]::PEAR1-VENUS* plant before (**e**) and after 24 hours of induction (**f**). *REV* exhibits a distinct transcriptional pattern where its expression is initially uniform in vascular tissue (**e'**), and highest expression is localized in PSE, while decreasing towards xylem axis (**e''** and **e'''**). When PEAR1-VENUS is overexpressed (**f**), the transcription pattern of *REV* is also activated in the central domain of vascular tissue, resulting in the radial symmetric *REV* transcription pattern. **g-h**, The expression pattern of *pATHB8::HTA6-YFP* is highly specific to xylem cells (**g**), and its expression is enhanced after 24 hours of induction of PEAR1 overexpression with a broad expression domain (**h**). Number in each panel indicates samples with similar results of the total independent biological samples analysed.

811

**Extended Data Fig. 10| The boundary between HD-ZIP III and PEAR proteins forms within the PSE-IN.**

**a**, Summarizing results on pattern of auxin-cytokinin (data shown in Extended Data Fig. 7a), HD-ZIP III (*CNA*, data shown in Extended Data Fig. 8g and k) and PEAR (PEAR1, data shown in Fig. 1d and Extended Data Fig. 4a-d) in procambium. In the simulation, we simulate the concentration of HD-ZIP III and PEAR1 along the axis between metaxylem (MX, brown arrowheads) and PSE (white arrowheads). **b**, Summarizing result of procambial development. Number of cells between metaxylem and PSE increases during procambial development (data shown in Extended Data Fig. 1). Therefore, the model is defined as a line in one spatial dimension representing 3, 4 or 5 cells from the centre of the xylem axis to the outer edge of the PSE cell. **c**, The regulatory network embedded within each cell. Regulatory interactions shown using a black line have been published previously, whilst those using a green line are described for the first time here. **d**, Predicted concentration gradient of all elements in 3, 4 or 5 cells (from left to right). Within different root geometries corresponding to different growth stages in *Arabidopsis*, both PEAR1 and HD-ZIP III are co-localized in PSE-IN, forming a sharp concentration boundary within this cell. In **f-h**, only PEAR and HD-ZIP concentrations are shown, whilst all model components are shown here in all cases. **e**, Quantification of expression level of *CNA-GFP* and PEAR1-GFP at 4-cell region in wild-type background. *n* represents individual measurements across 3 (*CNA-GFP*) or 4 (*PEAR1-GFP*) independent biological samples, respectively. Bar graphs represent mean. Error bars are s.d. Dots, individual data points. **f**, Including all three interactions labelled in panel **c** in the model, results in the formation of sharp concentration gradients of PEAR1 (black line) and HD-ZIP III (blue line) with the boundary forming in the PSE-IN. **g**, In simulations where HD-ZIP III does not regulate PEAR1 diffusion (Interaction (3) in panel **c**), PEAR1 protein is predicted to spread into the procambium and metaxylem as shown in Fig. 4k-m and Extended Data Fig. 4. **h**, In simulations where PEAR1 does not activate *HD-ZIP III* transcription (Interaction (1) in panel **c**), the concentration of HD-ZIP III is reduced in the PSE-IN cell as shown in Fig. 4h-j. **i**, A regulatory mechanism forming the boundary between a dividing and a non-dividing cell during

838 procambial development. White, orange, dark-green and light-green arrowheads indicate PSE, PSE-  
839 LN, PSE-IN and IPC. Scale bars, 25  $\mu$ m.

840  
841 **Supplementary Table 1|**  
842 This file contains a list of primers and plasmids used in this study.

843  
844 **Supplementary Table 2|**  
845 This file contains a list of putative PEAR1/PEAR2 direct targets and their description.

846  
847 **Supplementary Table 3|**  
848 This file contains individual *P*-values for Tukey's HSD test (for Fig. 2f and Extended Data Fig. 5b-d)  
849 and for two-sided Student's *t*-test (for Extended Data Fig. 7b). For Tukey's HSD test, data contains:  
850 difference in mean values (diff), the lower end point of the interval (lwr), the upper end point of the  
851 interval (upr) and *p*-value after adjustment for the multiple comparisons (*p* adj).  
852